arXiv:1611.04219v1 [math.CO] 14 Nov 2016

Resistance distance and Kirchhoff index in corona-vertex and corona-edge of subdivision graph

Qun Liu [a], Jia-Bao Liu [b,†], Shaohui Wang [c], Masoud Karimi [d,e]

a. School of Mathematics and Statistics, Hexi University, Gansu, Zhangye, 734000, P.R. China

b. School of Mathematic and Physics, Anhui Jianzhu University, Hefei, 230601, P.R. China

c. Department of Mathematic and Computer Science, Adelphi University, Garden City, NY 11550, USA

d. School of Mathematical Sciences, Anhui University, Hefei, 230601, P.R. China

e. Department of Mathematics, Bojnourd Branch, Islamic Azad University, Bojnourd, Iran

Abstract

The subdivision graph $S(G)$ of a graph G is the graph obtained by inserting a new vertex into every edge of G. In [11], two classes of new corona graphs, the corona-vertex of the subdivision graph $G_1 \diamondsuit G_2$ and corona-edge of the subdivision graph $G_1 \star G_2$ were defined. The adjacency spectrum and the signless Laplacian spectrum of the two new graphs were computed when G_1 is an arbitrary graph and G_2 is an r-regular graph. In this paper, we give the formulate of the resistance distance and the Kirchhoff index in $G_1 \diamondsuit G_2$ and $G_1 \star G_2$ when G_1 and G_2 are arbitrary graphs. These results generalize them in [11].

Keywords: Kirchhoff index, Resistance distance, Group inverse, Corona

AMS Mathematics Subject Classification(2000): 05C25, 05C50; 15A09

1 Introduction

Throughout this article, all graphs considered are simple and undirected. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G) = \{v_1, v_2, ..., v_n\}$ and edge set $E(G) = \{e_1, e_2, ..., e_m\}$. In 1993, Klein and Randić [1] introduced a distance function named resistance distance on the basis of electrical network theory. They view a graph as an electrical network and each edge of the graph is assumed to be a unit resistor, then take the resistance distance between vertices to be the effective resistance between them. Let G be a simple graph with the vertex set $V(G) = \{v_1, v_2, ..., v_n\}$, and $r_{ij}(G)$ denote the effective resistance distance between vertices v_i and v_j as computed with Ohm′s law when all the edges of G are considered to be unit

† Corresponding author. E-mail addresses: liuqun09@yeah.net, liujiabaoad@163.com, shaohui-wang@yahoo.com, karimimth@yahoo.com.

resistors. The sum of resistance distance $Kf(G) = \sum_{u,v \in V(G)} r_{uv}(G)$ was proposed in Ref. [1], later called the Kirchhoff index of G in Ref. [2]. The resistance distance and the Kirchhoff index attracted extensive attention due to its wide applications in physics, chemistry and others. For more information on resistance distance and Kirchhoff index of graphs, the readers are referred to Refs. [8-10, 14-19, 21-29] and the references therein.

Let $G = (V(G), E(G))$ be a graph with vertex set $V(G)$ and edge set $E(G)$. Let d_i be the degree of vertex i in G and $D_G = diag(d_1, d_2, \cdots, d_{|V(G)|})$ the diagonal matrix with all vertex degrees of G as its diagonal entries. For a graph G, let A_G and R_G denote the adjacency matrix and vertex-edge incidence matrix of G, respectively. The matrix $L_G = D_G - A_G$ is called the Laplacian matrix of G, where D_G is the diagonal matrix of vertex degrees of G. The line graph $l(G)$ of a graph G is the graph $V(l(G)) = E(G)$ such that two vertices in $l(G)$ are adjacent if and only if the corresponding edges in G are adjacent. Note that $R_G R_G^T = D_G + A_G$ and $R_G^T R_G = A(l(G)) + 2I_m$, where $m = |E(G)|$. The $\{1\}$-inverse of M is a matrix X such that $MXM = M$. If M is singular, then it has infinite $\{1\}$- inverse [4]. We use $M^{(1)}$ to denote any $\{1\}$- inverse of a matrix M, and let $(M)_{uv}$ denote the (u, v)- entry of M. It is known that resistance distances in a connected graph G can be obtained from any $\{1\}$- inverse of L_G ([3], [5]).

The subdivision graph $S(G)$ of a graph G is the graph obtained by inserting a new vertex into every edge of G. The set of such new vertices is denoted by $I(G)$. In Ref. [11], two new graph operations based on subdivision graphs: the corona-vertex and the corona-edge of the subdivision graph were introduced. The corona-vertex of the subdivision graph of two vertex-disjoint graphs G_1 and G_2 denoted by $G_1 \diamondsuit G_2$, is the graph obtained from G_1 and $|V(G_1)|$ copies of $S(G_2)$ by joining the ith vertex of $V(G_1)$ to every vertex in the ith copy of G_2. The corona-edge of the subdivision graph of two vertex-disjoint graphs G_1 and G_2, denoted by $G_1 \star G_2$, is the graph obtained from G_1 and $|V(G_1)|$ copies of $S(G_2)$ by joining the ith vertex of $V(G_1)$ to every vertex in the ith copy of $I(G_2)$.

Note that if G_1 is a graph on n_1 vertices and m_1 edges, and G_2 is a graph on n_2 vertices and m_2 edges, then the corona-vertex of the subdivision graph $G_1 \diamondsuit G_2$ has $n_1(1 + n_2 + m_2)$ vertices and $m_1 + n_1 n_2 + 2n_1 m_2$ edges, and the corona-edge of the subdivision graph $G_1 \star G_2$ has $n_1(1 + n_2 + m_2)$ vertices and $m_1 + 3n_1 m_2$ edges.

It is of interest to study the Kirchhoff index of graphs derived from a single graph. The formulas and lower bounds of the Kirchhoff index of the line graph, subdivision graph of a connected regular graph are reported in Refs. [5,18]. The formulas for the resistance distance and Kirchhoff index of the subdivision graphs of general graphs were obtained in Refs. ([26], [27]). In Ref. [7], Bu et al. obtained the formulae for resistance distances in subdivision-vertex join and subdivision-edge join. In [12], Liu gave the resistance distance and Kirchhoff index of R-vertex join and R-edge join of two graphs. Recently Liu et al [13] gave the $\{1\}$-inverse of the Laplacian of subdivision-vertex and subdivision-edge coronae. Motivated by these work, in this paper, we formulate the resistance distance and the Kirchhoff index in $G_1 \diamondsuit G_2$ and $G_1 \star G_2$ respectively.

2 Preliminaries

For a square matrix M, the group inverse of M, denoted by $M^{\#}$, is the unique matrix X such that $MXM = M$, $XMX = X$ and $MX = XM$. It is known that $M^{\#}$ exists if and only if $rank(M) = rank(M^2)$. If M is real symmetric, then $M^{\#}$ exists and $M^{\#}$ is a symmetric $\{1\}$-

inverse of M. Actually, $M^{\#}$ is equal to the Moore-Penrose inverse of M since M is symmetric.

We use $M^{(1)}$ to denote any $\{1\}$- inverse of a matrix M. Let $(M)_{uv}$ denote the (u,v) entry of M.

Lemma 2.1([3], [6]) Let G be a connected graph. Then

$$r_{uv}(G) = (L_G^{(1)})_{uu} + (L_G^{(1)})_{vv} - (L_G^{(1)})_{uv} - (L_G^{(1)})_{vu} = (L_G^{\#})_{uu} + (L_G^{\#})_{vv} - 2(L_G^{\#})_{uv}.$$

Let $\mathbf{1}_n$ denotes the column vector of dimension n with all the entries equal one. We will often use $\mathbf{1}$ to denote an all-ones column vector if the dimension can be read from the context.

Lemma 2.2 ([20]) For any graph, we have $L_G^{\#}\mathbf{1} = 0$.

Lemma 2.3 ([30]) Let

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

be a nonsingular matrix. If A and D are nonsingular, then

$$\begin{aligned} M^{-1} &= \begin{pmatrix} A^{-1} + A^{-1}BS^{-1}CA^{-1} & -A^{-1}BS^{-1} \\ -S^{-1}CA^{-1} & S^{-1} \end{pmatrix} \\ &= \begin{pmatrix} (A - BD^{-1}C)^{-1} & -A^{-1}BS^{-1} \\ -S^{-1}CA^{-1} & S^{-1} \end{pmatrix}, \end{aligned}$$

where $S = D - CA^{-1}B$.

Lemma 2.4 ([20]) Let

$$M = \begin{pmatrix} L_1 & L_2 \\ L_2^T & L_3 \end{pmatrix}$$

be a symmetric matrix and L_1 is nonsingular. Then

$$X = \begin{pmatrix} L_1^{-1} + L_1^{-1}L_2S^{\#}L_2^TL_1^{-1} & -L_1^{-1}L_2S^{\#} \\ -S^{\#}L_2^TL_1{-1} & S^{\#} \end{pmatrix}$$

is a symmetric $\{1\}$-inverse of M, where $S = L_3 - L_2^TL_1^{-1}L_2$.

For a vertex i of a graph G, let $T(i)$ denote the set of all neighbors of i in G.

Lemma 2.5([7]) Let G be a connected graph. For any $i, j \in V(G)$,

$$r_{ij}(G) = d_i^{-1}(1 + \sum_{k \in T(i)} r_{kj}(G) - d_i^{-1} \sum_{k,l \in T(i)} r_{kl}(G)).$$

Lemma 2.6([20]) Let G be a connected graph on n vertices. Then

$$Kf(G) = ntr(L_G^{(1)}) - \mathbf{1}^T L_G^{(1)}\mathbf{1} = ntr(L_G^{\#}).$$

Lemma 2.7 Let $\mathbf{1}_n$ and $j_{n\times m}$ be all-one column vector of dimensions n and all-one $n \times m$ matrix, respectively. For any matrix A, we have

$$(\mathbf{1}_{m_2} \otimes I_{n_1})A_{n_1\times n_1}(\mathbf{1}_{n_2}^T \otimes I_{n_1}) = j_{m_2\times n_2} \otimes A_{n_1\times n_1},$$

where $\otimes$ denote the Kronecker product of matrices $A = (a_{ij})$ and B, i.e $A \otimes B$ is defined to be the partition matrix $(a_{ij}B)$.

Proof.

$$\begin{aligned}(\mathbf{1}_{m_2} \otimes I_{n_1})A_{n_1\times n_1}(\mathbf{1}_{n_2}^T \otimes I_{n_1}) &= \begin{pmatrix} I_{n_1} \\ I_{n_1} \\ \vdots \\ I_{n_1} \end{pmatrix}_{(n_1m_2)\times n_1} A_{n_1\times n_1} \begin{pmatrix} I_{n_1} & I_{n_1} & \cdots & I_{n_1} \end{pmatrix}_{n_1\times(n_1n_2)} \\ &= \begin{pmatrix} A & A & \cdots & A \\ A & A & \cdots & A \\ \vdots & \vdots & \ddots & \vdots \\ A & A & \cdots & A \end{pmatrix}_{(n_1m_2)\times(n_1n_2)} \\ &= j_{m_2\times n_2} \otimes A_{n_1\times n_1}.\end{aligned}$$

□

3 Resistance distance in corona-vertex and corona-edge of subdivision graph

We first give formulae for resistance distance in $G_1 \diamondsuit G_2$.

Theorem 3.1 Let G_1 be a graph on n_1 vertices and m_1 edges and G_2 be a graph on n_2 vertices and m_2 edges. Then the following hold:

(i) For any $i, j \in V(G_1)$, we have

$$r_{ij}(G_1 \diamondsuit G_2) = (L^{\#}_{G_1})_{ii} + (L^{\#}_{G_1})_{jj} - 2(L^{\#}_{G_1})_{ij}.$$

(ii) For any $i, j \in V(G_2)$, we have

$$r_{ij}(G_1 \diamondsuit G_2) = 2((L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1})_{ii} + 2((L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1})_{jj}$$

$$-2((L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1})_{ij}.$$

(iii) For any $i \in V(G_1)$, $j \in V(G_2)$, we have

$$r_{ij}(G_1 \diamondsuit G_2) = (L^{\#}_{G_1})_{ii} + 2((L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1})_{jj} - (L^{\#}_{G_1})_{ij}.$$

(iv) For any $i \in I(G_2)$, $j \in V(G_1) \cup V(G_2)$, let $u_iv_i \in E(G_2)$ denote the edge corresponding to i, we have

$$r_{ij}(G_1 \diamondsuit G_2) = \frac{1}{2} + \frac{1}{2}r_{u_ij}(G_1 \diamondsuit G_2) + \frac{1}{2}r_{v_ij}(G_1 \diamondsuit G_2) - \frac{1}{4}r_{u_iv_i}(G_1 \diamondsuit G_2).$$

(v) For any $i, j \in I(G_2)$, let $u_iv_i, u_jv_j \in E(G_2)$ denote the edges corresponding to i, j, we have

$$\begin{aligned}r_{ij}(G_1 \diamondsuit G_2) &= \frac{1}{2} + \frac{1}{2}r_{u_ij}(G_1 \diamondsuit G_2) + \frac{1}{2}r_{v_ij}(G_1 \diamondsuit G_2) - \frac{1}{4}r_{u_iv_i}(G_1 \diamondsuit G_2) \\ &= 1 + \frac{1}{4}[r_{u_iu_j}(G_1 \diamondsuit G_2) + r_{u_iv_j}(G_1 \diamondsuit G_2) + r_{v_iu_j}(G_1 \diamondsuit G_2) \\ &\quad + r_{v_iv_j}(G_1 \diamondsuit G_2) - r_{u_iv_i}(G_1 \diamondsuit G_2) - r_{u_jv_j}(G_1 \diamondsuit G_2)].\end{aligned}$$

Proof. Let R_2 be the incidence matrix of G_2. Then with a proper labeling of vertices (see Ref.[11]), the Laplacian matrix of $G_1 \diamondsuit G_2$ can be written as

$$L(G_1 \diamondsuit G_2) = \begin{pmatrix} 2I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} & 0_{n_1 m_2 \times n_1} \\ -R_2 \otimes I_{n_1} & (D_{G_2} + I_{n_2}) \otimes I_{n_1} & -\mathbf{1}_{n_2} \otimes I_{n_1} \\ 0_{n_1 \times n_1 m_2} & -\mathbf{1}_{n_2}^T \otimes I_{n_1} & L_{G_1} + n_2 I_{n_1} \end{pmatrix},$$

where $0_{s,t}$ denotes the $s \times t$ matrix with all entries equal to zero. Let

$$L_1 = \begin{pmatrix} 2I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} \\ -R_2 \otimes I_{n_1} & (D_{G_2} + I_{n_2}) \otimes I_{n_1} \end{pmatrix},$$

$L_2 = \begin{pmatrix} 0_{n_1 m_2 \times n_1} \\ -\mathbf{1}_{n_2} \otimes I_{n_1} \end{pmatrix}$, $L_2^T = \begin{pmatrix} 0_{n_1 \times n_1 m_2} & -\mathbf{1}_{n_2}^T \otimes I_{n_1} \end{pmatrix}$, and $L_3 = L_{G_1} + n_2 I_{n_1}$.

We begin with the calculation about L_1^{-1}. By Lemma 2.3, we have

$$\begin{aligned} R_1 &= [(D_{G_2} + I_{n_2}) \otimes I_{n_1} - (R_2 \otimes I_{n_1})(2I_{m_2} \otimes I_{n_1})^{-1}(R_2^T \otimes I_{n_1})]^{-1} \\ &= [(D_{G_2} + I_{n_2}) \otimes I_{n_1} - \frac{1}{2} R_2 R_2^T \otimes I_{n_1}]^{-1} \\ &= [(D_{G_2} + I_{n_2}) - \frac{1}{2}(D_{G_2} + A_{G_2})]^{-1} \otimes I_{n_1} \\ &= 2(L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1}. \end{aligned}$$

One can obtain that

$$L_1^{-1} = \begin{pmatrix} T_1 & (R_2^T (L_{G_2} + 2I_{n_2})^{-1}) \otimes I_{n_1} \\ ((L_{G_2} + 2I_{n_2})^{-1} R_2) \otimes I_{n_1} & 2(L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1} \end{pmatrix},$$

where $T_1 = \frac{1}{2}(I_{m_2} + R_2^T (2I_{n_2} + L_{G_2})^{-1} R_2) \otimes I_{n_1}$.

Now we are ready to calculate S. Let

$$S = L_{G_1} + n_2 I_{n_1} - \begin{pmatrix} 0_{n_1 \times n_1 m_2} & -\mathbf{1}_{n_2}^T \otimes I_{n_1} \end{pmatrix} \begin{pmatrix} 2I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} \\ -R_2 \otimes I_{n_1} & A \end{pmatrix}^{-1} \begin{pmatrix} 0_{n_1 m_2 \times n_1} \\ -\mathbf{1}_{n_2} \otimes I_{n_1} \end{pmatrix},$$

where $A = (D_{G_2} + I_{n_2}) \otimes I_{n_1}$, then

$$\begin{aligned} S &= L_{G_1} + n_2 I_{n_1} - (\mathbf{1}_{n_2}^T ((D_{G_2} + I_{n_2}) I_{n_2} - \frac{1}{2} R_2 R_2^T)^{-1} \mathbf{1}_{n_2}) \otimes I_{n_1}) \\ &= L_{G_1} + n_2 I_{n_1} - (\mathbf{1}_{n_2}^T (\frac{1}{2} L_{G_2} + I_{n_2})^{-1} \mathbf{1}_{n_2}) \otimes I_{n_1} \\ &= L_{G_1}. \end{aligned}$$

By Lemma 2.4, we have $S^{\#} = L_{G_1}^{\#}$.

Next, according to Lemma 2.4, we calculate $-L_1^{-1} L_2 S^{\#}$ and $-L_1^{-1} L_2 S^{\#} L_2^T L_1^{-1}$.

$$\begin{aligned} -L_1^{-1} L_2 S^{\#} &= -\begin{pmatrix} T_1 & (R_2^T (L_{G_2} + 2I_{n_2})^{-1}) \otimes I_{n_1} \\ ((L_{G_2} + 2I_{n_2})^{-1} R_2) \otimes I_{n_1} & 2(L_{G_2} + 2I_{n_2})^{-1} \otimes I_{n_1} \end{pmatrix} \begin{pmatrix} 0_{n_1 m_2 \times n_1} \\ -\mathbf{1}_{n_2} \otimes I_{n_1} \end{pmatrix} S^{\#} \\ &= \begin{pmatrix} (R_2^T (L_{G_2} + 2I_{n_2})^{-1} \mathbf{1}_{n_2}) \otimes I_{n_1} \\ (2(L_{G_2} + 2I_{n_2})^{-1} \mathbf{1}_{n_2}) \otimes I_{n_1} \end{pmatrix} S^{\#}. \end{aligned}$$

Note that $(L_{G_2}+2I_{n_2})\mathbf{1}_{n_2}=2\cdot\mathbf{1}_{n_2}$, then

$$R_2^T(L_{G_2}+2I_{n_2})^{-1}\mathbf{1}_{n_2}=\frac{1}{2}R_2^T\mathbf{1}_{n_2}=\mathbf{1}_{m_2},\qquad 2(L_{G_2}+2I_{n_2})^{-1}\mathbf{1}_{n_2}=\mathbf{1}_{n_2},$$

For convenience, let $H=\mathbf{1}_{m_2}\otimes I_{n_1}, K=\mathbf{1}_{n_2}\otimes I_{n_1}$, then

$$\begin{aligned}-L_1^{-1}L_2S^{\#} &= \begin{pmatrix}\mathbf{1}_{m_2}\otimes I_{n_1}\\ \mathbf{1}_{n_2}\otimes I_{n_1}\end{pmatrix}S^{\#}\\ &= \begin{pmatrix}HS^{\#}\\ KS^{\#}\end{pmatrix},\end{aligned}$$

and let $Q=(L_{G_2}+2I_{n_2})$, then

$$\begin{aligned}-L_1^{-1}L_2S^{\#}L_2^TL_1^{-1} &= \begin{pmatrix}-HS^{\#}\\ -KS^{\#}\end{pmatrix}\begin{pmatrix}0_{n_1\times n_1m_2} & -K^T\end{pmatrix}\begin{pmatrix}T_1 & (R_2^TQ^{-1})\otimes I_{n_1}\\ (Q^{-1}R_2)\otimes I_{n_1} & 2Q^{-1}\otimes I_{n_1}\end{pmatrix}\\ &= \begin{pmatrix}-HS^{\#}\\ -KS^{\#}\end{pmatrix}\begin{pmatrix}(-\mathbf{1}_{n_2}^TQ^{-1}R_2)\otimes I_{n_1} & -(2\cdot\mathbf{1}_{n_2}^TQ^{-1}\otimes I_{n_1})\end{pmatrix}.\end{aligned}$$

Note that $\mathbf{1}_{n_2}^T(L_{G_2}+2I_{n_1})=2\cdot\mathbf{1}_{n_2}^T$, then $\mathbf{1}_{n_2}^TQ^{-1}R_2=\mathbf{1}_{n_2}^T(L_{G_2}+2I_{n_1})^{-1}R_2=\frac{1}{2}\mathbf{1}_{n_2}^TR_2=\mathbf{1}_{m_2}^T$, so

$$\begin{aligned}-L_1^{-1}L_2S^{\#}L_2^TL_1^{-1} &= \begin{pmatrix}-HS^{\#}\\ -KS^{\#}\end{pmatrix}\begin{pmatrix}-\mathbf{1}_{m_2}^T\otimes I_{n_1} & -\mathbf{1}_{n_2}^T\otimes I_{n_1}\end{pmatrix}\\ &= \begin{pmatrix}HS^{\#}H^T & HS^{\#}K^T\\ KS^{\#}H^T & KS^{\#}K^T\end{pmatrix}.\end{aligned}$$

Based on Lemma 2.3 and 2.4, the following matrix

$$N=\begin{pmatrix}T_1+HS^{\#}H^T & R_2^TQ^{-1}\otimes I_{n_1}+HS^{\#}K^T & HS^{\#}\\ Q^{-1}R_2\otimes I_{n_1}+KS^{\#}H^T & 2Q^{-1}\otimes I_{n_1}+KS^{\#}K^T & KS^{\#}\\ S^{\#}H & S^{\#}K & S^{\#}\end{pmatrix}\tag{3.1}$$

is a symmetric $\{1\}$- inverse of $L(G_1\diamondsuit G_2)$, where $T_1=\frac{1}{2}(I_{m_2}+R_2^T(2I_{n_2}+L_{G_2})^{-1}R_2)\otimes I_{n_1}$, $Q=(L_{G_2}+2I_{n_2})$, $H=\mathbf{1}_{m_2}\otimes I_{n_1}, K=\mathbf{1}_{n_2}\otimes I_{n_1}$.

For any $i,j\in V(G_1)$, by Lemma 2.1 and the Equation (3.1), we have

$$r_{ij}(G_1\diamondsuit G_2) = (L_{G_1}^{\#})_{ii}+(L_{G_1}^{\#})_{jj}-2(L_{G_1}^{\#})_{ij}.$$

For any $i,j\in V(G_2)$, by Lemma 2.1 and the Equation (3.1), we have

$$r_{ij}(G_1\diamondsuit G_2)=2((L_{G_2}+2I_{n_2})^{-1}\otimes I_{n_1})_{ii}+2((L_{G_2}+2I_{n_2})^{-1}\otimes I_{n_1})_{jj}$$

$$-2((L_{G_2}+2I_{n_2})^{-1}\otimes I_{n_1})_{ij}.$$

For any $i\in V(G_1)$, $j\in V(G_2)$, by Lemma 2.1 and the Equation (3.1), we have

$$r_{ij}(G_1\diamondsuit G_2) = (L_{G_1}^{\#})_{ii}+2((L_{G_2}+2I_{n_2})^{-1}\otimes I_{n_1})_{jj}-(L_{G_1}^{\#})_{ij}.$$

For any $i\in I(G_2)$, $j\in V(G_1)\cup V(G_2)$, Let $u_iv_i\in E(G_2)$ denote the edge corresponding to i, By Lemma 2.5, we have

$$r_{ij}(G_1\diamondsuit G_2) = \frac{1}{2}+\frac{1}{2}r_{u_ij}(G_1\diamondsuit G_2)+\frac{1}{2}r_{v_ij}(G_1\diamondsuit G_2)-\frac{1}{4}r_{u_iv_i}(G_1\diamondsuit G_2).$$

For any $i,j \in I(G_2)$, let $u_iv_i, u_jv_j \in E(G_2)$ denote the edges corresponding to i,j. By Lemma 2.5, we have

$$\begin{aligned} r_{ij}(G_1 \diamondsuit G_2) &= \frac{1}{2} + \frac{1}{2}r_{u_ij}(G_1 \diamondsuit G_2) + \frac{1}{2}r_{v_ij}(G_1 \diamondsuit G_2) - \frac{1}{4}r_{u_iv_i}(G_1 \diamondsuit G_2) \\ &= 1 + \frac{1}{4}[r_{u_iu_j}(G_1 \diamondsuit G_2) + r_{u_iv_j}(G_1 \diamondsuit G_2) + r_{v_iu_j}(G_1 \diamondsuit G_2) \\ &\quad + r_{v_iv_j}(G_1 \diamondsuit G_2) - r_{u_iv_i}(G_1 \diamondsuit G_2) - r_{u_jv_j}(G_1 \diamondsuit G_2)]. \end{aligned}$$

□

When G_2 is a regular graph, we will give the formulae for resistance distance in $G_1 \star G_2$ as follows.

Theorem 3.2 Let G_1 be an arbitrary graph on n_1 vertices, and G_2 an r_2- regular graph on n_2 vertices and m_2 edges, then the following hold:

(i) For any $i,j \in V(G_1)$, we have

$$r_{ij}(G_1 \star G_2) = (L^{\#}_{G_1})_{ii} + (L^{\#}_{G_1})_{jj} - 2(L^{\#}_{G_1})_{ij}.$$

(ii) For any $i,j \in V(G_2)$, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= (3(r_2I_{n_2} + L_{G_2}) \otimes I_{n_1})^{-1}_{ii} + (3(r_2I_{n_2} + L_{G_2}) \otimes I_{n_1})^{-1}_{jj} \\ &\quad - 2(3(r_2I_{n_2} + L_{G_2}) \otimes I_{n_2})^{-1}_{ij}. \end{aligned}$$

(iii) For any $i \in V(G_1)$, $j \in V(G_2)$, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= (L^{\#}_{G_1})_{ii} + (3(r_2I_{n_2} + L(G_2) \otimes I_{n_1})^{-1}_{jj} \\ &\quad - (L^{\#}_{G_1})_{ij}. \end{aligned}$$

(iv) For any $i \in I(G_2)$, $j \in V(G_1) \cup V(G_2)$, let $u_iv_i \in E(G_2)$ denote the edge corresponding to i, $k \in V(G_1)$, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= \frac{1}{3}(1 + r_{u_ij}(G_1 \star G_2) + r_{v_ij}(G_1 \star G_2) + r_{kj}(G_1 \star G_2)) \\ &\quad - \frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_ik}(G_1 \star G_2) + r_{v_ik}(G_1 \star G_2)). \end{aligned}$$

(v) For any $i,j \in I(G_2)$, let $u_iv_i, u_jv_j \in E(G_2)$ denote the edges corresponding to i,j, $k \in V(G_1)$, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= \frac{1}{3}(1 + r_{ju_i}(G_1 \star G_2) + r_{jv_i}(G_1 \star G_2) + r_{jk}(G_1 \star G_2) \\ &\quad - \frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_ik}(G_1 \star G_2) + r_{v_ik}(G_1 \star G_2)) \\ &= \frac{1}{3}(1 + r_{iu_j}(G_1 \star G_2) + r_{iv_j}(G_1 \star G_2) + r_{ik}(G_1 \star G_2) \\ &\quad - \frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_jk}(G_1 \star G_2) + r_{v_jk}(G_1 \star G_2)). \end{aligned}$$

Proof. Let R_2 be the incidence matrix of G_2. Then with a proper labeling of vertices (see Ref.[13]), the Laplacian matrix of $G_1 \star G_2$ can be written as

$$L(G_1 \star G_2) = \begin{pmatrix} 3I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} & -\mathbf{1}_{m_2} \otimes I_{n_1} \\ -R_2 \otimes I_{n_1} & r_2 I_{n_2} \otimes I_{n_1} & 0_{n_1 n_2 \times n_1} \\ -\mathbf{1}_{m_2}^T \otimes I_{n_1} & 0_{n_1 \times n_1 n_2} & L_{G_1} + m_2 I_{n_1} \end{pmatrix},$$

where $0_{s,t}$ denotes the $s \times t$ matrix with all entries equal to zero. Let

$$L_1 = \begin{pmatrix} 3I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} \\ -R_2 \otimes I_{n_1} & r_2 I_{n_2} \otimes I_{n_1} \end{pmatrix},$$

$L_2 = \begin{pmatrix} -\mathbf{1}_{m_2} \otimes I_{n_1} \\ 0_{n_1 n_2 \times n_1} \end{pmatrix}$, $L_2^T = \begin{pmatrix} -\mathbf{1}_{m_2}^T \otimes I_{n_1} & 0_{n_1 \times n_1 n_2} \end{pmatrix}$, and $L_3 = L_{G_1} + m_2 I_{n_1}$.

We begin with the calculation about L_1^{-1}. By Lemma 2.3, we have

$$\begin{aligned} R_2 &= [r_2 I_{n_2} \otimes I_{n_1} - (R_2 \otimes I_{n_1})(3I_{m_2} \otimes I_{n_1})^{-1}(R_2^T \otimes I_{n_1})]^{-1} \\ &= [r_2 I_{n_2} \otimes I_{n_1} - \frac{1}{3} R_2 R_2^T \otimes I_{n_1}]^{-1} \\ &= [r_2 I_{n_2} - \frac{1}{3}(r_2 I_{n_2} + A_{G_2})]^{-1} \otimes I_{n_1} \\ &= 3(L_{G_2} + r_2 I_{n_2})^{-1} \otimes I_{n_1}. \end{aligned}$$

One can obtain that

$$L_1^{-1} = \begin{pmatrix} T_2 & (R_2^T (L_{G_2} + r_2 I_{n_2})^{-1}) \otimes I_{n_1} \\ ((L_{G_2} + r_2 I_{n_2})^{-1} R_2) \otimes I_{n_1} & 3(L_{G_2} + r_2 I_{n_2})^{-1} \otimes I_{n_1} \end{pmatrix},$$

where $T_2 = \frac{1}{3}(I_{m_2} + R_2^T (r_2 I_{n_2} + L_{G_2})^{-1} R_2) \otimes I_{n_1}$.

Now we are ready to calculate S.

Let

$$S = L_{G_1} + m_2 I_{n_1} - \begin{pmatrix} -\mathbf{1}_{m_2}^T \otimes I_{n_1} & 0_{n_1 \times n_1 n_2} \end{pmatrix} \begin{pmatrix} 3I_{m_2} \otimes I_{n_1} & -R_2^T \otimes I_{n_1} \\ -R_2 \otimes I_{n_1} & r_2 I_{n_2} \otimes I_{n_1} \end{pmatrix}^{-1} \begin{pmatrix} -\mathbf{1}_{m_2} \otimes I_{n_1} \\ 0_{n_1 n_2 \times n_1} \end{pmatrix},$$

then

$$\begin{aligned} S &= L_{G_1} + m_2 I_{n_1} - (-\mathbf{1}_{m_2}^T \otimes I_{n_1})((3I_{m_2} - \frac{1}{r_2} R_2^T R_2)^{-1} \otimes I_{n_1})(-\mathbf{1}_{m_2} \otimes I_{n_1}) \\ &= L_{G_1} + m_2 I_{n_1} - (r_2 \mathbf{1}_{m_2}^T (L_{l(G_2)} + r_2 I_{m_2})^{-1} \mathbf{1}_{m_2}) \otimes I_{n_1} \\ &= L_{G_1}. \end{aligned}$$

By Lemma 2.4, we have $S^{\#} = L_{G_1}^{\#}$.

Similiarly, according to Lemma 2.4, we calculate $-L_1^{-1} L_2 S^{\#}$ and $-L_1^{-1} L_2 S^{\#} L_2^T L_1^{-1}$.

$$\begin{aligned} -L_1^{-1} L_2 S^{\#} &= \begin{pmatrix} T_2 & (R_2^T (L_{G_2} + r_2 I_{n_2})^{-1}) \otimes I_{n_1} \\ ((L_{G_2} + r_2 I_{n_2})^{-1} R_2) \otimes I_{n_1} & 3(L_{G_2} + r_2 I_{n_2})^{-1} \otimes I_{n_1} \end{pmatrix} \begin{pmatrix} -\mathbf{1}_{m_2} \otimes I_{n_1} \\ 0_{n_1 n_2 \times n_1} \end{pmatrix} S^{\#} \\ &= \begin{pmatrix} -T_2(\mathbf{1}_{m_2} \otimes I_{n_1}) \\ -((L_{G_2} + r_2 I_{n_2})^{-1} R_2) \otimes I_{n_1})(\mathbf{1}_{m_2} \otimes I_{n_1}) \end{pmatrix} S^{\#}. \end{aligned}$$

Note that $(L_{G_2}+r_2I_{n_2})\mathbf{1}_{n_2}=r_2\mathbf{1}_{n_2}$, then

$$\begin{aligned} T_2(1_{m_2}\otimes I_{n_1}) &= \frac{1}{3}((I_{m_2}+R_2^T(r_2I_{n_2}+L_{G_2})^{-1}R_2)\otimes I_{n_1})(\mathbf{1}_{m_2}\otimes I_{n_1}) \\ &= \frac{1}{3}(\mathbf{1}_{m_2}+R_2^T(r_2I_{n_2}+L_{G_2})^{-1}R_2\mathbf{1}_{m_2})\otimes I_{n_1} \\ &= \frac{1}{3}(\mathbf{1}_{m_2}+R_2^T(r_2I_{n_2}+L_{G_2})^{-1}r_2\mathbf{1}_{n_2})\otimes I_{n_1} \\ &= \frac{1}{3}(\mathbf{1}_{m_2}+R_2^T\mathbf{1}_{n_2})\otimes I_{n_1}=\mathbf{1}_{m_2}\otimes I_{n_1}. \end{aligned}$$

and $((r_2I_{n_2}+L_{G_2})^{-1}R_2)\otimes I_{n_1})(\mathbf{1}_{m_2}\otimes I_{n_1})=((r_2I_{n_2}+L_{G_2})^{-1}r_2\mathbf{1}_{n_2})\otimes I_{n_1}=\mathbf{1}_{n_2}\otimes I_{n_1}$, so

$$-L_1^{-1}L_2S^{\#} = \begin{pmatrix} -(\mathbf{1}_{m_2}\otimes I_{n_1}) \\ -(\mathbf{1}_{n_2}\otimes I_{n_1}) \end{pmatrix} S^{\#} = \begin{pmatrix} -(\mathbf{1}_{m_2}\otimes I_{n_1})S^{\#} \\ -(\mathbf{1}_{n_2}\otimes I_{n_1})S^{\#} \end{pmatrix}.$$

Let $C=L_{G_2}+r_2I_{n_2}$, $H=\mathbf{1}_{m_2}\otimes I_{n_1}, K=\mathbf{1}_{n_2}\otimes I_{n_1}$,

$$\begin{aligned} -L_1^{-1}L_2S^{\#}L_2^TL_1^{-1} &= \begin{pmatrix} -HS^{\#} \\ -KS^{\#} \end{pmatrix}\begin{pmatrix} -H^T & 0_{n_1\times n_1n_2} \end{pmatrix}\begin{pmatrix} T_2 & R_2^TC^{-1}\otimes I_{n_1} \\ C^{-1}R_2\otimes I_{n_1} & 3C^{-1}\otimes I_{n_1} \end{pmatrix} \\ &= \begin{pmatrix} -HS^{\#} \\ -KS^{\#} \end{pmatrix}\begin{pmatrix} -H^TT_2 & -H^T(R_2^TC^{-1}\otimes I_{n_1}) \end{pmatrix}. \end{aligned}$$

Note that $H^TT_2=(\mathbf{1}_{m_2}\otimes I_{n_1})^TT_2=(T_2(\mathbf{1}_{m_2}\otimes I_{n_1}))^T=H^T$, then

$$-L_1^{-1}L_2S^{\#}L_2^TL_1^{-1} = \begin{pmatrix} -HS^{\#} \\ -KS^{\#} \end{pmatrix}\begin{pmatrix} -H^T & K^T \end{pmatrix} = \begin{pmatrix} HS^{\#}H^T & HS^{\#}K^T \\ KS^{\#}H^T & KS^{\#}K^T \end{pmatrix}.$$

Based on Lemma 2.3 and 2.4, the following matrix

$$N=\begin{pmatrix} T_2+HS^{\#}H^T & R_2^TC^{-1}\otimes I_{n_1}+HS^{\#}K^T & HS^{\#} \\ C^{-1}R_2\otimes I_{n_1}+KS^{\#}H^T & 3C^{-1}\otimes I_{n_1}+KS^{\#}K^T & KS^{\#} \\ S^{\#}H & S^{\#}K & S^{\#} \end{pmatrix} \tag{3.2}$$

is a symmetric $\{1\}$- inverse of $L(G_1\star G_2)$, where $T_2=\frac{1}{3}(I_{m_2}+R_2^T(r_2I_{n_2}+L_{G_2})^{-1}R_2)\otimes I_{n_1}$, $C=(L_{G_2}+r_2I_{n_2})$, $H=\mathbf{1}_{m_2}\otimes I_{n_1}, K=\mathbf{1}_{n_2}\otimes I_{n_1}$.

For any $i,j\in V(G_1)$, by Lemma 2.1 and the Equation (3.2), we have

$$r_{ij}(G_1\star G_2) = (L_{G_1}^{\#})_{ii}+(L_{G_1}^{\#})_{jj}-2(L_{G_1}^{\#})_{ij}.$$

For any $i,j\in V(G_2)$, by Lemma 2.1 and the Equation (3.2), we have

$$\begin{aligned} r_{ij}(G_1\star G_2) &= (3(r_2I_{n_2}+L_{G_2})\otimes I_{n_1})_{ii}^{-1}+(3(r_2I_{n_2}+L_{G_2})\otimes I_{n_1})_{jj}^{-1} \\ &\quad -2(3(r_2I_{n_2}+L_{G_2})\otimes I_{n_1})_{ij}^{-1}. \end{aligned}$$

For any $i\in V(G_1)$, $j\in V(G_2)$, by Lemma 2.1 and the Equation (3.2), we have

$$\begin{aligned} r_{ij}(G_1\star G_2) &= (L_{G_1}^{\#})_{ii}+(3(r_2I_{n_2}+L(G_2)\otimes I_{n_1})_{jj}^{-1} \\ &\quad -(L_{G_1}^{\#})_{ij}. \end{aligned}$$

For any $i \in I(G_2)$, $j \in V(G_1) \cup V(G_2)$, let $u_iv_i \in E(G_2)$ denote the edge corresponding to i, $k \in V(G_1)$, by Lemma 2.5, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= \frac{1}{3}(1 + r_{u_ij}(G_1 \star G_2) + r_{v_ij}(G_1 \star G_2) + r_{kj}(G_1 \star G_2)) \\ &\quad -\frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_ik}(G_1 \star G_2) + r_{v_ik}(G_1 \star G_2)). \end{aligned}$$

For any $i, j \in I(G_2)$, let $u_iv_i, u_jv_j \in E(G_2)$ denote the edges corresponding to i, j, $k \in V(G_1)$, by Lemma 2.5, we have

$$\begin{aligned} r_{ij}(G_1 \star G_2) &= \frac{1}{3}(1 + r_{ju_i}(G_1 \star G_2) + r_{jv_i}(G_1 \star G_2) + r_{jk}(G_1 \star G_2) \\ &\quad -\frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_ik}(G_1 \star G_2) + r_{v_ik}(G_1 \star G_2)) \\ &= \frac{1}{3}(1 + r_{iu_j}(G_1 \star G_2) + r_{iv_j}(G_1 \star G_2) + r_{ik}(G_1 \star G_2) \\ &\quad -\frac{1}{9}(r_{u_iv_i}(G_1 \star G_2) + r_{u_jk}(G_1 \star G_2) + r_{v_jk}(G_1 \star G_2)). \end{aligned}$$

□

4 Kirchhoff index in corona-vertex and corona-edge of subdivision graph

Theorem 4.1 Let G_1 be a graph on n_1 vertices and m_1 edges and G_2 be a graph on n_2 vertices and m_2 edges. Then

$$\begin{aligned} Kf(G_1 \diamondsuit G_2) &= n_1(1 + n_2 + m_2)[\frac{n_1m_2}{2} + \frac{n_1}{2}(tr(Q^{-1}A_{G_2}) + tr(Q^{-1}D_{G_2})) + 2n_1\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} \\ &\quad + \frac{m_2 + n_2 + 1}{n_1}Kf(G_1)] - \frac{n_1}{2}(\pi^TQ^{-1}\pi) - \frac{5n_1m_2 + 2n_1n_2}{2}, \end{aligned}$$

where $Q = L_{G_2} + 2I_{n_2}$, $D_{G_2} = diag(d_1, d_2, \cdots, d_{n_2})$ and $\pi = (d_1, d_2, \cdots, d_{n_2})^T$.

Proof. Let $L^{(1)}_{G_1 \diamondsuit G_2}$ be the symmetric $\{1\}$-inverse of $L_{G_1 \diamondsuit G_2}$. Then

$$tr(L^{(1)}_{G_1 \diamondsuit G_2}) = tr(T_1 + HS^{\#}H^T) + tr(2Q^{-1} \otimes I_{n_1} + KS^{\#}K^T) + tr(S^{\#}),$$

where $T_1 = \frac{1}{2}(I_{m_2} + R_2^T(2I_{n_2} + L_{G_2})^{-1}R_2) \otimes I_{n_1}$, $Q = (L_{G_2} + 2I_{n_2})$, $S^{\#} = L^{\#}_{G_1}$, $H = \mathbf{1}_{m_2} \otimes I_{n_1}$, $K = \mathbf{1}_{n_2} \otimes I_{n_1}$, and R_2 is the incidence matrix of G_2.

By Lemma 2.7, $HS^{\#}H^T = j_{m_2 \times m_2} \otimes S^{\#}$, $KS^{\#}K^T = j_{n_2 \times n_2} \otimes S^{\#}$, then

$$tr(HS^{\#}H^T) = m_2tr(S^{\#}), \quad tr(KS^{\#}K^T) = n_2tr(S^{\#}).$$

Note that the eigenvalues of Q are $\mu_1(G_2) + 2, ..., \mu_n(G_2) + 2$, then

$$tr(Q^{-1}) = \sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2) + 2}.$$

Recall that $R_2R_2^T = D_{G_2} + A_{G_2}$, then

$$tr(R_2^TQ^{-1}R_2) = tr(Q^{-1}R_2R_2^T) = tr(Q^{-1}A_{G_2}) + tr(Q^{-1}D_{G_2}).$$

Thus, substituting $tr(R_2^TQ^{-1}R_2)$, $tr(Q^{-1})$ into $tr(L^{(1)}_{G_1\diamondsuit G_2})$, we have

$$\begin{aligned}
tr(L^{(1)}_{G_1\diamondsuit G_2}) &= tr(T_1) + tr(j_{m_2\times m_2}\otimes S^{\#}) + 2tr(Q^{-1}\otimes I_{n_1}) + tr(j_{n_2\times n_2}\otimes S^{\#}) + tr(S^{\#}) \\
&= \frac{n_1m_2}{2} + \frac{n_1}{2}tr(R_2^TQ^{-1}R_2) + m_2tr(S^{\#}) + 2n_1tr(Q^{-1}) + n_2tr(S^{\#}) + tr(S^{\#}) \\
&= \frac{n_1m_2}{2} + \frac{n_1}{2}(tr(Q^{-1}A_{G_2}) + tr(Q^{-1}D_{G_2})) + m_2tr(S^{\#}) \\
&\quad +2n_1\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} + n_2tr(S^{\#}) + tr(S^{\#}) \\
&= \frac{n_1m_2}{2} + \frac{n_1}{2}(tr(Q^{-1}A_{G_2}) + tr(Q^{-1}D_{G_2})) + 2n_1\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} \\
&\quad +\frac{m_2+n_2+1}{n_1}Kf(G_1).
\end{aligned}$$

Next, we calculate the $\mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1}$. Since $L_G^{\#}\mathbf{1} = 0$, by Lemma 2.7, then

$$\mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1} = \mathbf{1}^TT_1\mathbf{1} + \mathbf{1}^T(2Q^{-1}\otimes I_{n_1})\mathbf{1} + \mathbf{1}^T((R_2^TQ^{-1})\otimes I_{n_1})\mathbf{1} + \mathbf{1}^T((Q^{-1}R_2^T)\otimes I_{n_1})\mathbf{1}.$$

Note that $R_2\mathbf{1} = \pi$, where $\pi = (d_1, d_2, \cdots, d_{n_2})^T$, then

$$\mathbf{1}^T((R_2^TQ^{-1}R_2)\otimes I_{n_1})\mathbf{1} = n_1(\pi^TQ^{-1}\pi).$$

Since $Q\mathbf{1}_{n_2} = 2\cdot\mathbf{1}_{n_2}$, then $\mathbf{1}^T(2Q^{-1}\otimes I_{n_1})\mathbf{1} = 2n_1(\mathbf{1}^TQ^{-1}\mathbf{1}) = n_1n_2$ and

$$\mathbf{1}^T((Q^{-1}R_2)\otimes I_{n_1})\mathbf{1} = n_1\mathbf{1}^T(Q^{-1}R_2)\mathbf{1} = \frac{1}{2}\mathbf{1}^TR_2\mathbf{1} = \frac{1}{2}n_1\sum_{i=1}^{n_2}d_i = n_1m_2,$$

Similarly, $\mathbf{1}^T((R_2^TQ^{-1})\otimes I_{n_1})\mathbf{1} = (\mathbf{1}^T((Q^{-1}R_2)\otimes I_{n_1})\mathbf{1})^T = n_1m_2$.

Substituting $\mathbf{1}^T((R_2^TQ^{-1}R_2)\otimes I_{n_1})\mathbf{1}$, $\mathbf{1}^T((Q^{-1}R_2)\otimes I_{n_1})\mathbf{1}$, $\mathbf{1}^T((R_2^TQ^{-1})\otimes I_{n_1})\mathbf{1}$ into $\mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1}$, we get

$$\mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1} = \frac{n_1m_2}{2} + \frac{n_1}{2}(\pi^TQ^{-1}\pi) + 2m_2n_1 + n_1n_2.$$

Lemma 2.6 implies that

$$Kf(L^{(1)}_{G_1\diamondsuit G_2}) = n_1(1+n_2+m_2)tr(L^{(1)}_{G_1\diamondsuit G_2}) - \mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1}.$$

Then plugging $tr(L^{(1)}_{G_1\diamondsuit G_2})$ and $\mathbf{1}^T(L^{(1)}_{G_1\diamondsuit G_2})\mathbf{1}$ into the equation above, we obtain the required result. □

Corollary 4.2 Let G_1 be a graph on n_1 vertices and m_1 edges and G_2 an r_2- regular graph on n_2 vertices and m_2 edges. Then

$$\begin{aligned}
Kf(G_1\diamondsuit G_2) &= n_1(1+n_2+m_2)[\frac{n_1m_2}{2} + \frac{n_1}{2}(2r_2\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} - \sum_{i=1}^{n_2}\frac{\mu_i(G_2)}{\mu_i(G_2)+2}) + \\
&\quad 2n_1\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} + \frac{m_2+n_2+1}{n_1}Kf(G_1)] - \frac{n_1n_2r_2^2}{4} - \frac{5n_1m_2+2n_1n_2}{2}.
\end{aligned}$$

Proof. Since $tr(Q^{-1}D_{G_2}) = r_2tr(Q^{-1}) = r_2\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2}$, $tr(Q^{-1}A_{G_2}) = r_2tr(Q^{-1}) - tr(Q^{-1}L_{G_2}) = r_2\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+2} - \sum_{i=1}^{n_2}\frac{\mu_i(G_2)}{\mu_i(G_2)+2}$, $\pi^TQ^{-1}\pi = r^2\mathbf{1}^TQ^{-1}\mathbf{1} = \frac{n_2r_2^2}{2}$, then the required result is obtained by plugging $tr(Q^{-1}D_{G_2})$, $tr(Q^{-1}A_{G_2})$ and $\pi^TQ^{-1}\pi$ into Theorem 4.1. □

Theorem 4.3 Let G_1 be an arbitrary graph on n_1 vertices, and G_2 an r_2- regular graph on n_2 vertices and m_2 edges. Then

$$\begin{aligned} Kf(G_1 \star G_2) &= n_1(1+n_2+m_2)[\frac{n_1m_2}{3} + \frac{n_1}{3}(tr(C^{-1}A_{G_2}) + r_2\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+r_2}) + \\ & 3n_1\sum_{i=1}^{n_2}\frac{1}{\mu_i(G_2)+r_2} + \frac{m_2+n_2+1}{n_1}Kf(G_1)] - \frac{n_1m_2r_2 + n_1n_2(r_2+3)^2}{3r_2}, \end{aligned}$$

where $C = L_{G_2} + r_2I_{n_2}$.

Proof. The proof is similar to that of Theorem 4.1 and hence we omit details. □

Acknowledgment: This work was supported by the National Natural Science Foundation of China (Nos. 11461020, 11561042, 11601006 and 11471016), the Youth Foundation of Hexi University in Gansu Province (No.QN2013-07) and the Teaching Reform Project of Hexi University (No. HXXXJY-2014-011).

References

[1] D.J. Klein, M. Randić, Resistance distance. Journal of Mathematical Chemistry. 12(1), (1993) 81-95 .

[2] D. Bonchev, A.T. Balaban, X. Liu, D.J. Klein, Molecular cyclicity and centricity of polycyclic graphs. I: cyclicity based on resistance distances or reciprocal distances. International journal of quantum chemistry, 50(1), (1994) 1-20 .

[3] R.B. Bapat, S. Gupta, Resistance distance in wheels and fans. Indian Journal of Pure and Applied Mathematics. 41(1), (2010) 1-13 .

[4] A. Ben-Israel, T.N.E. Greville, Generalized inverses: Theory and Applications", Second ed., Springer-Verlag, New York, 2003.

[5] R.B. Bapat, Graphs and matrices. Universitext, Springer/Hindustan Book Agency, London/ New Delhi, 2010.

[6] C. Bu, L. Sun, J. Zhou, Y. Wei, A note on block representations of the group inverse of Laplacian matrices. Electronic Journal of Linear Algebra. 23(1), (2012) 866-876 .

[7] C. Bu, B. Yan, X. Zhou, J. Zhou, Resistance distance in subdivision-vertex join and subdivision-edge join of graphs. Linear Algebra and its Applications. 458(1), (2014) 454-462.

[8] H.Y. Chen, F.J. Zhang, Resistance distance and the normalized Laplacian spectrum. Discrete Applied Mathematics. 155(5), (2007) 654-661 .

[9] X. Gao, Y. Luo, W. Liu, Kirchhoff index in line, subdivision and total graphs of a regular graph. Discrete Applied Mathematics. 160(4), (2012) 560-565 .

[10] D.J. Klein, Resistance-distance sum rules. Croatica Chemica Acta. 75(2), (2002) 633-649 .

[11] P.L. Lu, Y. F. Miao, A-Spectra and Q-Spectra of Two Classes of Corona Graphs, Journal of Donghua University (English Edition). 3(1), (2014) 224-228 .

[12] X. Liu, J. Zhou, C. Bu, Resistance distance and Kirchhoff index of R-vertex join and R-edge join of two graphs. Discrete Applied Mathematics. 187(1), (2015) 130-139.

[13] J.B. Liu, X.F. Pan, Fu-Tao Hu, The {1}-inverse of the Laplacian of subdivision-vertex and subdivision-edge coronae with applications, Linear and Multilinear Algebra(http://dx.doi.org/10.1080/03081087.2016.1179249).

[14] J.B. Liu, X.F. Pan, L. Yu, D. Li, Complete characterization of bicyclic graphs with minimal Kirchhoffindex, Discret. Appl. Math. 200 (2016) 95-107.

[15] J.B. Liu, W.R. Wang, Y.M. Zhang, X.F. Pan, On degree resistance distance of cacti, Discret. Appl. Math. 203 (2016) 217-225.

[16] J.B. Liu, X.F. Pan, F.T. Hu, F.F. Hu, Asymptotic Laplacian-energy-like invariant of lattices. Appl. Math. Comput. 253, (2015) 205-214.

[17] J.B. Liu, J. Cao, The resistance distances of electrical networks based on Laplacian generalized inverse. Neurocomputing. 167, (2015) 306-313 .

[18] J.B. Liu, X.F. Pan, A unified approach to the asymptotic topological indices of various lattices. Appl. Math. Comput. 270, (2015) 62-73 .

[19] Q. Liu, J.B. Liu, J. Cao, The Laplacian polynomial and Kirchhoff index of graphs based on R-graphs, Neurocomputing. (2015), in press.

[20] L.Z. Sun, W.Z. Wang, J. Zhou, C. Bu, Some results on resistance distances and resistance matrices. Linear and Multilinear Algebra. 63(3), (2015) 523-533 .

[21] W. Wang, D. Yang, Y. Luo, The Laplacian polynomial and Kirchhoff index of graphs derived from regular graphs. Discrete Applied Mathematics, 161(18), (2013) 3063-3071 .

[22] S. Wang, B. Wei, Multiplicative Zagreb indices of k-trees, *Discrete Applied Mathematics 180 (2015) 168-175.*

[23] C. Wang, S. Wang, B. Wei, Cacti with Extremal PI Index, *Transactions on Combinatorics 5 (2016) 1-8.*

[24] W. J. Xiao, I.Gutman, Resistance distance and Laplacian spectrum, Theoretical Chemistry Accounts. 110(4) (2003) 284-289.

[25] Y.J. Yang, D.J. Klein, A recursion formula for resistance distances and its applications, Discrete Applied Mathematics, 161(16)(2013) 2702-2715 .

[26] Y.J. Yang, Kirchhoff index of subdivisions of graphs, Discrete Applied Mathematics. 171(1), (2014) 153-157.

[27] Y.J. Yang, D.J. Klein, Resistance distance-based graph invariants of subdivisions and triangulations of graphs, Discrete Applied Mathematics. 181(1) (2015) 260-274 .

[28] H.P. Zhang, Y.J. Yang, Resistance distance and Kirchhoff index in circulant graphs, International journal of quantum chemistry. 107(2) (2007) 330-339 .

[29] B. Zhou, N. Trinajstić , On resistance-distance and Kirchhoff index, Journal of mathematical chemistry, 46(1) 283-289 (2009).

[30] J. Zhou, L. Sun, W. Wang, C. Bu, Line star sets for Laplacian eigenvalues, Linear Algebra and its Applications, 440(1) 164-176 (2014).